

Size of the neutralino–nucleon cross–section in the light of a new determination of the pion–nucleon sigma term

A. Bottino^a, F. Donato^b, N. Fornengo^a, S. Scopel^{a *}

^a *Dipartimento di Fisica Teorica, Università di Torino
and INFN, Sez. di Torino, Via P. Giuria 1, I-10125 Torino, Italy*

^b *Laboratoire de Physique Théorique LAPTH
Annecy-le-Vieux, 74941, France*

Abstract

We discuss the implications of a new determination of the pion–nucleon sigma term for the evaluation of the neutralino–nucleon cross–section, and, in turn, for detection rates for relic neutralinos in WIMP direct experiments and in some of the indirect searches. It is shown that the new range for the pion–nucleon sigma term, taken at its face value, favours values of the neutralino–nucleon cross–sections which are sizeably larger than some of the current estimates. Implications for neutralino cosmological properties are derived.

I. INTRODUCTION

The evaluation of detection rates for relic neutralinos in WIMP direct experiments and in measurements of signals due to possible neutralino–neutralino annihilation in Earth and Sun [1] requires a reliable calculation of the neutralino–nucleon cross–section.

In Ref. [2] we stressed that sizeable uncertainties in the neutralino–nucleon cross–section are due to the estimate of the hadronic matrix elements, which describe the quark density

*E–mail: bottino@to.infn.it, donato@lapp.in2p3.fr, fornengo@to.infn.it,
scopel@to.infn.it

contents in the nucleon; this point was also remarked in Refs. [3,4]. In [2] we quantified the range of these uncertainties and showed that, within their ranges, the couplings of neutralinos to nucleon may be sizeably stronger than some of those currently employed in the literature. Subsequently, an analysis of this problem was also undertaken in Refs. [5–8].

One of the crucial ingredients in the calculation of the hadronic matrix elements is the so-called pion–nucleon sigma term. Its numerical derivation from experimental data of pion–nucleon scattering is rather involved, and thus the origin of considerable uncertainties. Recent experimental results in pion–nucleon scattering have now prompted a new determination of the pion–nucleon sigma term [9]. In the present paper, we consider the implications of these new inputs for the neutralino–nucleon cross-section; in particular, we show that the new data reinforce our previous conclusions of Ref. [2]. As compared to Ref. [2], the present paper, apart from the use of the new evaluation for the sigma term, contains also some other relevant updatings, mainly: i) radiatively corrected Higgs–quark couplings [10] and ii) recent experimental bounds on Higgs masses and supersymmetric parameters from LEP2 [11] and CDF [12].

II. NEUTRALINO–NUCLEON ELASTIC CROSS-SECTION

The neutralino–nucleon scalar cross-section may be written as

$$\sigma_{\text{scalar}}^{(\text{nucleon})} = \frac{8G_F^2}{\pi} M_Z^2 m_{\text{red}}^2 \left[\frac{F_h I_h}{m_h^2} + \frac{F_H I_H}{m_H^2} + \frac{M_Z}{2} \sum_q \langle N | \bar{q} q | N \rangle \sum_i P_{\tilde{q}_i} (A_{\tilde{q}_i}^2 - B_{\tilde{q}_i}^2) \right]^2. \quad (1)$$

The first two terms inside the brackets refer to the diagrams with exchanges of the two CP-even neutral Higgs bosons, h and H , in the t-channel (the exchange diagram of the CP-odd one, A , is strongly kinematically suppressed and then omitted here) [13] and the third term refers to the graphs with squark-exchanges in the s- and u-channels [14]. The mass m_{red} is the neutralino–nucleon reduced mass. Since, for simplicity, in the present paper we explicitly discuss only the Higgs-mediated terms, we do not report the expressions for the squark propagator $P_{\tilde{q}_i}$ and for the couplings $A_{\tilde{q}_i}$, $B_{\tilde{q}_i}$, which may be found in Ref. [15]. However, in the numerical results reported in this paper also the squark-exchange terms are included. The quark matrix elements $\langle N | \bar{q} q | N \rangle$ are meant over the nucleonic state.

The quantities $F_{h,H}$ and $I_{h,H}$ are defined as follows

$$\begin{aligned} F_h &= (-a_1 \sin \theta_W + a_2 \cos \theta_W)(a_3 \sin \alpha + a_4 \cos \alpha) \\ F_H &= (-a_1 \sin \theta_W + a_2 \cos \theta_W)(a_3 \cos \alpha - a_4 \sin \alpha) \\ I_{h,H} &= \sum_q k_q^{h,H} m_q \langle N | \bar{q} q | N \rangle, \end{aligned} \quad (2)$$

where the a_i 's are the coefficients in the definition of the neutralino as the lowest-mass linear superposition of photino ($\tilde{\gamma}$), zino ($\tilde{Z}$) and the two higgsino states ($\tilde{H}_1^\circ$, $\tilde{H}_2^\circ$)

$$\chi \equiv a_1 \tilde{\gamma} + a_2 \tilde{Z} + a_3 \tilde{H}_1^\circ + a_4 \tilde{H}_2^\circ. \quad (3)$$

The coefficients $k_q^{h,H}$ for up-type and down-type quarks are given in Table 1, in terms of the angle β , defined as $\tan \beta = \langle H_2^0 \rangle / \langle H_1^0 \rangle$ and the angle α , which rotates $H_1^{(0)}$ and $H_2^{(0)}$ into h and H . In Table I also included are the coefficients $k_q^{h,H}$ for the CP-odd Higgs boson A , which are important for the evaluation of the neutralino relic abundance. The entries include those radiative corrections which may be sizeable at large $\tan \beta$. These corrections affect the couplings to down-type quarks k_{d-type} , and are parametrized in terms of the quantity $\epsilon \equiv 1/(1 + \Delta)$, where Δ enters in the relationship between the fermion running masses m_d and the corresponding Yukawa couplings h_d [10]:

$$m_d = h_d \langle H_0^1 \rangle (1 + \Delta) \quad (4)$$

These corrections take contributions mainly from gluino-squark, chargino-squark and neutralino-stau loops [10].

III. EVALUATION OF THE QUANTITIES $M_Q \langle N | \bar{Q} Q | N \rangle$

For the calculation of the quantities $m_q \langle N | \bar{q} q | N \rangle$ it is useful to express them in terms of the pion-nucleon sigma term

$$\sigma_{\pi N} = \frac{1}{2} (m_u + m_d) \langle N | \bar{u} u + \bar{d} d | N \rangle, \quad (5)$$

of the quantity σ_0 , related to the size of the SU(3) symmetry breaking, and defined as

$$\sigma_0 \equiv \frac{1}{2} (m_u + m_d) \langle N | \bar{u} u + \bar{d} d - 2 \bar{s} s | N \rangle, \quad (6)$$

and of the ratio $r = 2m_s/(m_u + m_d)$.

Assuming isospin invariance for quarks u and d , the quantities $m_q \langle N | \bar{q} q | N \rangle$ for light quarks may be written as

$$m_u \langle N | \bar{u} u | N \rangle \simeq m_d \langle N | \bar{d} d | N \rangle \simeq \frac{1}{2} \sigma_{\pi N} \quad (7)$$

$$m_s \langle N | \bar{s} s | N \rangle \simeq \frac{1}{2} r (\sigma_{\pi N} - \sigma_0). \quad (8)$$

For the heavy quarks c, b, t , use of the heavy quark expansion [16] provides

$$\begin{aligned} m_c < N|\bar{c}c|N > &\simeq m_b < N|\bar{b}b|N > \simeq m_t < N|\bar{t}t|N > \simeq \\ &\simeq \frac{2}{27} \left[m_N - \sigma_{\pi N} + \frac{1}{2}r(\sigma_{\pi N} - \sigma_0) \right], \end{aligned} \quad (9)$$

where m_N is the nucleon mass. The quantities $I_{h,H}$ can then be rewritten as

$$I_{h,H} = k_{u\text{-type}}^{h,H} g_u + k_{d\text{-type}}^{h,H} g_d, \quad (10)$$

where

$$\begin{aligned} g_u &\simeq m_l < N|\bar{l}l|N > + 2 m_h < N|\bar{h}h|N > \\ &\simeq \frac{4}{27} (m_N + \frac{19}{8}\sigma_{\pi N} - \frac{1}{2}r(\sigma_{\pi N} - \sigma_0)), \end{aligned} \quad (11)$$

$$\begin{aligned} g_d &\simeq m_l < N|\bar{l}l|N > + m_s < N|\bar{s}s|N > + m_h < N|\bar{h}h|N > \\ &\simeq \frac{2}{27} (m_N + \frac{23}{4}\sigma_{\pi N} + \frac{25}{4}r(\sigma_{\pi N} - \sigma_0)); \end{aligned} \quad (12)$$

l stands for light quarks ($l = u, d$) and h denotes the heavy ones ($h = c, b, t$).

We recall that from the values of $\sigma_{\pi N}$ and σ_0 one can derive the fractional strange-quark content of the nucleon y

$$y = 2 \frac{< N|\bar{s}s|N >}{< N|\bar{u}u + \bar{d}d|N >}, \quad (13)$$

using the expression

$$y = 1 - \frac{\sigma_0}{\sigma_{\pi N}}. \quad (14)$$

A. Values for $\sigma_{\pi N}$, σ_0 and r

The range of $\sigma_{\pi N}$ we used previously in Ref. [2] is

$$41 \text{ MeV} \lesssim \sigma_{\pi N} \lesssim 57 \text{ MeV}. \quad (15)$$

This was derived from the pion–nucleon scattering amplitude, calculated at the so–called Cheng–Dashen point by Koch [17], and the evolution of the nucleon scalar form factor, as a function of the momentum transfer from $t = 2m_\pi^2$ to $t = 0$, evaluated in Ref. [18].

We now consider the new determination of $\sigma_{\pi N}$ presented in Ref. [9]. The George Washington University/TRIUMF group, using up–dated pion–nucleon scattering data [20] and a new partial–wave and dispersion relation analysis program, has derived a range for $\sigma_{\pi N}$ [9]

$$55 \text{ MeV} \lesssim \sigma_{\pi N} \lesssim 73 \text{ MeV}, \quad (16)$$

which turns out to be sizeably larger than the one of Eq. (15). Values of the nucleon scalar form factor at the Cheng–Dashen point higher than those of Ref. [17] were also reported in Ref. [21].

In the present paper we consider the effect of employing the new range for $\sigma_{\pi N}$, given in Eq. (16), in the evaluation of the quantities $m_q \langle N | \bar{q}q | N \rangle$'s.

Here, σ_0 is taken in the range [19]

$$\sigma_0 = 30 \div 40 \text{ MeV}. \quad (17)$$

Also the mass ratio $r = 2m_s/(m_u + m_d)$ may be affected by significant uncertainties [2]. However, here, for consistency with some of the previous determinations, we use the *default* value $r = 25$.

It has to be noted that combining together values of $\sigma_{\pi N}$ and σ_0 within their ranges in Eqs. (16) and (17) leads to rather large values for the fractional strange–quark content of the nucleon, as given by Eq. (14). This is a puzzle that urges further investigation in hadron physics. In what follows, in the variations of the quantities $\sigma_{\pi N}$ and σ_0 we impose the constraint that, anyway, $y \leq 0.5$.

B. Values for $m_q \langle N | \bar{q}q | N \rangle$, g_u and g_d

Inserting the numerical values of $\sigma_{\pi N}$, σ_0 and r in the expressions given at the beginning of this section, we finally obtain estimates for the quantities $m_q \langle N | \bar{q}q | N \rangle$, g_u and g_d . Since $m_s < N | \bar{s}s | N >$ is the most important term among the $m_q < N | \bar{q}q | N >$'s [22], unless $\tan \beta$ is very small, the extremes of the range of the neutralino–nucleon cross–section are provided by the extremes of the range for $m_s < N | \bar{s}s | N >$. These, in turn, are given by: $(m_s < N | \bar{s}s | N >)^{min} \simeq \frac{1}{2}r(\sigma_{\pi N}^{min} - \sigma_0^{max})$ and $(m_s < N | \bar{s}s | N >)^{max} \simeq \frac{1}{4}r\sigma_{\pi N}^{max}$ (to satisfy the constraint $y \leq 0.5$). We call *Set b* the one with $m_s < N | \bar{s}s | N > = (m_s < N | \bar{s}s | N >)^{min}$ and

Set c the one with $m_s < N|\bar{s}s|N\rangle = (m_s < N|\bar{s}s|N\rangle)^{max}$; we denote by *Set a* the reference set of Ref. [4]. The values of the quantities $m_q\langle N|\bar{q}q|N\rangle$, g_u and g_d are given in Table II.

We note that, with the new values of $\sigma_{\pi N}$, the coefficient g_d , which usually dominates in the neutralino–nucleon cross–section, turns out to fall in the range

$$266 \text{ MeV} \lesssim g_d \lesssim 523 \text{ MeV}, \quad (18)$$

to be compared with the reference value $g_d = 241 \text{ MeV}$ of *Set a*. Thus, for most supersymmetric configurations, one expects $(\sigma_{scalar}^{(nucleon)})_{Set b}/(\sigma_{scalar}^{(nucleon)})_{Set a} \simeq 1.5$ and $(\sigma_{scalar}^{(nucleon)})_{Set c}/(\sigma_{scalar}^{(nucleon)})_{Set a} \simeq 6$.

IV. RESULTS AND CONCLUSIONS

The supersymmetric model employed in the present paper for the calculation of the neutralino–nucleon cross–section and for the neutralino relic abundance is the one defined in Ref. [23,24] and denoted there as effMSSM. We refer to [23,24] for details on the theoretical aspects and on the updated experimental bounds.

Inserting the values of Table II into Eq.(1), one obtains the results displayed in Fig. 1. The two ratios in the cross–sections are plotted as a function of $\xi\sigma_{scalar}^{(nucleon)}$, where ξ is a rescaling factor for the neutralino local density. ξ is taken to be $\xi = \min\{1, \Omega_\chi h^2/(\Omega_m h^2)_{min}\}$, in order to have rescaling, when $\Omega_\chi h^2$ turns out to be less than $(\Omega_m h^2)_{min}$ (here $(\Omega_m h^2)_{min}$ is set to the value 0.05).

The use of $\xi\sigma_{scalar}^{(nucleon)}$ instead of simply $\sigma_{scalar}^{(nucleon)}$ allows one to better identify the range of sensitivity in current WIMP direct experiments [25], in Fig. 1. Taking into account astrophysical uncertainties [26], this range may be established to be

$$4 \cdot 10^{-10} \text{ nbarn} \lesssim \xi\sigma_{scalar}^{(nucleon)} \lesssim 2 \cdot 10^{-8} \text{ nbarn}, \quad (19)$$

for WIMP masses in the interval $40 \text{ GeV} \lesssim m_W \lesssim 200 \text{ GeV}$. In Fig. 1 we notice that, in the sensitivity range of Eq.(19), the cross–section ratios are actually of the sizes obtained by our previous estimate based on dominance of the term g_d , *i. e.* $(\sigma_{scalar}^{(nucleon)})_{Set b}/(\sigma_{scalar}^{(nucleon)})_{Set a} \simeq 1.5$ and $(\sigma_{scalar}^{(nucleon)})_{Set c}/(\sigma_{scalar}^{(nucleon)})_{Set a} \simeq 6$.

We turn now to the implications that our present analysis has for the cosmological properties of relic neutralinos, as explored by present WIMP direct detection experiments. Fig. 2 provides the essential information. The detection of relic neutralinos would be of great

relevance even if these particles constitute only a subdominant dark matter population [24]; however, it is obvious that the most attractive case is when the neutralino relic abundance falls into the interval of cosmological interest: $0.05 \lesssim \Omega_m h^2 \lesssim 0.3$. Fig. 2 shows to what extent the most interesting region: $0.05 \lesssim \Omega_m h^2 \lesssim 0.3$ and $4 \cdot 10^{-10} \text{ nbarn} \lesssim \xi \sigma_{\text{scalar}}^{(\text{nucleon})} \lesssim 2 \cdot 10^{-8} \text{ nbarn}$, is covered by neutralino configurations, depending on the values of the input parameters of Table II. The up-right frontier of the scatter plots moves progressively upward, as we move from *Set a* of inputs to *Set b* and to *Set c*.

The main results of the present paper may be summarized as follows:

- The new range of the pion–nucleon sigma term favours values of the neutralino–nucleon cross-section which are sizeably larger (a factor of 1.5 to 6) than some of the current estimates. However, a word of caution has to be said here: the new derivation of the pion–nucleon sigma term implies a rather high value for the fractional strange-quark content of the nucleon; this point requires further investigation in the framework of hadron physics. Thus, for instance, the consistency between the present indication for higher values of the pion–nucleon sigma term and the determination of the range of σ_0 has to be understood.
- Uncertainties implied by the hadronic quantities for the neutralino–nucleon cross-section are still quite sizeable; these have to be taken in due consideration in evaluations of the neutralino event rates.
- The previous remark applies not only to event rates for WIMP direct detection, but also to the evaluation of the neutrino fluxes expected at neutrino telescopes, as a consequence of possible neutralino–neutralino annihilations inside the Earth and the Sun. In fact, these fluxes depend on the capture rate of neutralinos by the macroscopic bodies; in turn, this rate depends on the neutralino–nucleon cross-section.
- A larger size of the neutralino–nucleon cross-section, as indicated by the new hadronic data, implies that present WIMP direct experiments (and some indirect measurements) explore larger sectors of the supersymmetric space where the neutralino may be of cosmological interest.

REFERENCES

- [1] For a review of neutralino as a dark matter particle, see, for instance, A. Bottino and N. Fornengo, *Proceedings of the Fifth School on Non-Accelerator Particle Astrophysics* (Abdus Salam International Centre for Theoretical Physics, Trieste), Eds. R.A. Carrigan, Jr., G. Giacomelli and N. Paver, E.U.T. 1999, [arXiv:hep-ph/9904469].
- [2] A. Bottino, F. Donato, N. Fornengo and S. Scopel, *Astroparticle Physics* **13** (2000) 215, [arXiv:hep-ph/9909228].
- [3] A. Bottino, V. de Alfaro, N. Fornengo, A. Morales, J. Puimedon, S. Scopel, *Mod.Phys.Lett.* **A7** (1992) 733.
- [4] G. Jungman, M. Kamionkowski, K. Griest, *Phys. Rep.* **267** (1996) 195.
- [5] J. Ellis, A. Ferstl and K.A. Olive, *Phys. Lett.* **B481** (2000) 304, [arXiv:hep-ph/0001005].
- [6] E. Accomando, R. Arnowitt, B. Dutta and Y. Santoso, *Nucl.Phys.* **B585** (2000) 124, [arXiv:hep-ph/0001019].
- [7] A. Corsetti and P. Nath, [arXiv:hep-ph/0003186].
- [8] J.L. Feng, K.T. Matchev and F. Wilczek, *Phys.Lett.* **B482** (2000) 388-399, [arXiv:hep-ph/0004043].
- [9] M.M. Pavan, R.A. Arndt, I.I. Strakovsky and R.L. Workman, [arXiv:hep-ph/0111066].
- [10] M. Carena, S. Mrenna and E. E. M. Wagner, *Phys. Rev.* **D60** (1999) 075010; H. Eberl, K. Hidaka, S. Kraml, W. Majerotto and Y. Yamada, *Phys. Rev.* **D62** (2000) 055006; A. Djouadi and M. Drees, *Phys. Lett.* **B 484** (2000) 183.
- [11] I.M. Fisk and K. Nagai, talks at the XXXth Int. Conf. on High Energy Physics, Osaka, July 2000, <http://www.ichep2000.hep.sci.osaka-u.ac.jp>; P.J. Donan (ALEPH Collaboration), March 2000, http://alephwww.cern.ch/ALPUB/seminar/lepc_mar2000/lepc2000.pdf.
- [12] J.A.Valls (CDF Coll.) FERMILAB-Conf-99/263-E CDF; <http://fnalpubs.fnal.gov/archive/1999/conf/Conf-99-263-E.html>.
- [13] R. Barbieri, M. Frigeni and G.F. Giudice, *Nucl. Phys.* **B313** (1989) 725.
- [14] K. Griest, *Phys. Rev.* **D38** (1988) 2357, *Nucl. Phys.* **B313** (1989) 725.
- [15] A. Bottino, F. Donato, N. Fornengo and S. Scopel, *Phys. Rev* **D59** (1999)095003 [arXiv:hep-ph/9808456].
- [16] M.A. Shifman, A.I. Vainshtein and V.I. Zacharov, *Phys. Lett.* **B78** (1978) 443; *JEPT Lett.* **22** (1975) 55.
- [17] R. Koch, *Z. Phys.***C15** (1982) 161.
- [18] J. Gasser, H. Leutwyler and M.E. Sainio, *Phys. Lett.* **B253** (1991) 260.
- [19] J. Gasser, H. Leutwyler, *Phys. Rep.***87** (1982) 77.
- [20] SAID pion-nucleon database, <http://gwdac.phys.gwu.edu/>.
- [21] M.G. Olsson, [arXiv:hep-ph/0001203].
- [22] G.B. Gelmini, P. Gondolo and E. Roulet, *Nucl. Phys.* **B351** (1991) 623.
- [23] A. Bottino, F. Donato, N. Fornengo and S. Scopel, *Phys. Rev* **D63** (2001) 125003 [arXiv:hep-ph/0010203].
- [24] A. Bottino, F. Donato, N. Fornengo and S. Scopel, [arXiv:hep-ph/0105233], to appear in the Proceedings of the 15th Rencontres de Physique de la Vallée d'Aoste: *Results and Perspective in Particle Physics*, La Thuile, March 2001.

- [25] E. Garcia et al., Phys. Rev. **D51** (1995) 1458; R. Bernabei et al., Phys. Lett. **B389** (1996) 757, Phys. Lett. **B 480** (2000) 23, Eur. Phys. J. **C 18** (2000) 283; Phys. Lett. **B 424** (1998) 195, Phys. Lett. **B436** (1998) 379; L. Baudis et al., Phys. Rev. **D59** (1999) 022001; A. Morales et al., Phys. Lett. **B489** (2000) 268 and [arXiv:hep-ex/0101037]; N.J.C. Spooner et al., Phys. Lett. **B473** (2000) 330; R. Abusaidi *et al.*, *Phys. Rev. Lett.* **84** (2000) 5699; A. Benoit et al., [arXiv:astro-ph/0106094].
- [26] P. Belli, R. Bernabei, A. Bottino, F. Donato, N. Fornengo, D. Prospero and S. Scopel, *Phys. Rev.* **D61** (1999) 023512, and references quoted therein.

TABLES

TABLE I. Values of the coefficients $k_q^{h,H}$ in Eq.(2).

	h	H	A
k_{u-type}	$\cos \alpha / \sin \beta$	$\sin \alpha / \sin \beta$	$1 / \tan \beta$
k_{d-type}	$-\sin \alpha / \cos \beta - \epsilon \cos(\alpha - \beta) \tan \beta$	$\cos \alpha / \cos \beta - \epsilon \sin(\alpha - \beta) \tan \beta$	$\tan \beta(1 + \epsilon)$

TABLE II. Values of the quantities $m_q \langle N | \bar{q} q | N \rangle$, g_u and g_d . *Set a* is the reference set given in Ref. [4]. *Set b* and *Set c* are the sets corresponding to the minimal and maximal values of $m_s < N | \bar{s} s | N \rangle$, respectively. All quantities are in units of MeV.

	$m_l < N \bar{q}_l q_l N \rangle$	$m_s < N \bar{s} s N \rangle$	$m_h < N \bar{h} h N \rangle$	g_u	g_d
<i>Set a</i> [4]	27	131	56	139	214
<i>Set b</i>	28	186	52	132	266
<i>Set c</i>	37	456	30	97	523

FIGURES




FIG. 1. Ratios of the neutralino–nucleon cross-sections, for values of the quantities $m_q \langle N | \bar{q}q | N \rangle$, g_u and g_d , as given in Table II




FIG. 2. Scatter plot of $\sigma_{\text{scalar}}^{(\text{nucleon})}$ versus $\Omega_\chi h^2$, when *Set a* is employed for the quantities given in Table II. The curve denoted by *Set a* visualizes the up-right frontier of the scatter plot. The curves denoted by *Set b* and by *Set c* give the locations of the frontiers of the scatter plots (not shown in this figure) obtained by using values of *Set b* and *Set c*, respectively. m_χ is taken in the range $40 \text{ GeV} \leq m_\chi \leq 200 \text{ GeV}$. The two horizontal lines bracket the sensitivity region defined by Eq. (19), when the rescaling factor ξ is set equal to one. The two vertical lines denote the range $0.05 \leq \Omega_m h^2 \leq 0.3$. Dots denote gauginos ($P > 0.9$), circles denote higgsinos ($P < 0.1$) and crosses denote mixed ($0.1 \leq P \leq 0.9$) configurations (P being defined as $P \equiv a_1^2 + a_2^2$).